arXiv:1407.3066v1 [math.NT] 11 Jul 2014

# An elementary proof of a power series identity for the weighted sum of all finite abelian $p$-groups

Pritam Majumder

*Indian Institute of Technology Kanpur, India*

**Abstract**

Using combinatorial techniques, we prove that the weighted sum of the inverse number of automorphisms of all finite abelian $p$-groups $\sum_G |G|^{-u} |\mathrm{Aut}(G)|^{-1}$ is equal to $\prod_{j=u+1}^{\infty} (1-1/p^j)^{-1}$, where $u$ is a non-negative integer. This result was originally obtained by H. Cohen and H. W. Lenstra, Jr. In this paper we give a new elementary proof of their result.

## 1. Introduction

Let $\mathcal{G}_p$ denote the set of all finite abelian $p$-groups, for a prime number $p$. We will give an elementary combinatorial proof of the following theorem:

**Theorem.** *For a prime $p$ and a non-negative integer $u$, the following holds*

$$\sum_{G \in \mathcal{G}_p} \frac{1}{|G|^u \cdot |\mathrm{Aut}(G)|} = \prod_{j=u+1}^{\infty} \left(1 - \frac{1}{p^j}\right)^{-1}.$$

The above result was obtained by Cohen and Lenstra in their famous paper [1]. Their approach is more complicated but has the advantage that it generalizes naturally to finite modules over the rings of integers of number fields. In the special case of $u = 0$, a nice combinatorial proof was given by Hall, which can be found in [2]. Our proof is a generalization of the proof of Hall.

## 2. Proof of the Theorem

For $m \geq 0$, let $a_m$ be the number of partitions of $m$ with each part at least $u+1$ and for $i, j \geq 0$, let $b_{i,j}$ be the number of partitions of $i$ with greatest part exactly equal to $j$.

**Lemma 1.** *For all $m \geq 0$, the following holds*

$$a_m = \sum_{i+uj=m} b_{i,j}.$$

*Proof.* We will give a bijection argument. Note that, the number of partitions of $i$ with greatest part $j$, where $i+uj = m$, is equal to the number of partitions of $i + uj = m$ with greatest part $j$ occurring at least $u+1$ times. Hence $\sum_{i+uj=m} b_{i,j}$ is equal to the number of partitions of $m$ with greatest part occurring at least $u+1$ times. Now, a partition of $m$ has greatest part occurring at least $u+1$ times if and only if it's conjugate partition has each part at least $u+1$. This gives a bijection between the partitions of $m$ with greatest part occurring at least $u+1$ times and the partitions of $m$ with each part at least $u+1$. Therefore, $\sum_{i+uj=m} b_{i,j} = a_m$. □

**Lemma 2.** *For each $n \geq 0$, let us define*

$$f_n(q) := \sum_{N=0}^{\infty} b_{N,n} q^N,$$

*which is a formal power series in $q$. Then,*

$$\prod_{j=u+1}^{\infty} (1-q^j)^{-1} = \sum_{n=0}^{\infty} f_n(q) q^{nu}.$$

*Proof.* Note that,

$$\prod_{j=u+1}^{\infty} (1-q^j)^{-1} = \sum_{m=0}^{\infty} a_m q^m$$

since, for each $m$, the coefficient of $q^m$ on LHS is equal to the number of partitions of $m$ with each part at least $u+1$. Then,

$$\begin{aligned}\sum_{n=0}^{\infty} f_n(q)q^{nu} &= \sum_{n=0}^{\infty}\left(\sum_{N=0}^{\infty} b_{N,n}q^N\right)q^{nu}\\ &= \sum_{n=0}^{\infty}\sum_{N=0}^{\infty} b_{N,n}q^{N+nu}\\ &= \sum_{m=0}^{\infty}\left(\sum_{i+uj=m} b_{i,j}\right)q^m\\ &= \sum_{m=0}^{\infty} a_m q^m\\ &= \prod_{j=u+1}^{\infty}(1-q^j)^{-1}.\end{aligned}$$

□

We also need the following lemma, which computes the cardinality of $\mathrm{Aut}(G)$, for a finite abelian $p$-group $G$.

**Lemma 3.** *Fix a prime $p$. Suppose $G$ is a finite abelian $p$-group and*

$$G = \prod_{i=1}^{k}(\mathbb{Z}/p^{e_i}\mathbb{Z})^{r_i}$$

*for some $k \geq 0$, $e_1 > e_2 > \cdots > e_k > 0$ and $r_i \geq 0$. Then*

$$|\mathrm{Aut}(G)| = \left(\prod_{i=1}^{k}\left(\prod_{s=1}^{r_i}(1-p^{-s})\right)\right)\left(\prod_{1\leq i,j\leq k} p^{\min(e_i,e_j)r_i r_j}\right).$$

*Proof.* See [3], Theorem 1.2.10. □

Now let us return to the proof of the theorem. We follow a similar argument given in [2] or in the proof of Theorem 2.1.2 of [3]. First, note that, there is an associated partition corresponding to every finite abelian $p$-group and corresponding to every partition there is an associated finite abelian $p$-group; this comes from writing finite abelian $p$-groups uniquely as a product

of cyclic groups. For example, if we write a finite abelian $p$-group $G$ as,

$$G = \prod_{i=1}^{k} \mathbb{Z}/p^{e_i}\mathbb{Z}$$

where $e_1 \geq e_2 \geq \cdots \geq e_k > 0$, then the associated partition $\lambda$ is given by $\lambda = (e_1, e_2, \ldots, e_k)$. And, corresponding to every partition $\lambda = (\lambda_1, \lambda_2, \ldots, \lambda_k)$ the associated $p$-group $G_\lambda$ is given by $G_\lambda = \prod_{i=1}^{k} \mathbb{Z}/p^{\lambda_i}\mathbb{Z}$. Note that, if $|\lambda|$ denotes the size of the partition $\lambda$, then the order of the $p$-group $G_\lambda$ is given by $|G_\lambda| = p^{|\lambda|}$.

Let $\lambda := (\lambda_1, \ldots, \lambda_l)$ be a partition of size $n$ and suppose $\lambda' := (\lambda'_1, \ldots, \lambda'_m)$ is it's conjugate partition. Then, note that, in $G_\lambda$ (as a product of cyclic groups), the factor $\mathbb{Z}/p^i\mathbb{Z}$ occurs exactly $\lambda'_i - \lambda'_{i+1}$ times (where $\lambda'_{m+1} := 0$). Then using Lemma 3 we can write

$$\begin{aligned}
|\mathrm{Aut}(G_\lambda)| &= \left(\prod_{i=1}^{m}\left(\prod_{s=1}^{\lambda'_i-\lambda'_{i+1}}(1-p^{-s})\right)\right)\left(\prod_{1\leq i,j\leq m} p^{\min(i,j)(\lambda'_i-\lambda'_{i+1})(\lambda'_j-\lambda'_{j+1})}\right) \\
&= \left(\prod_{i=1}^{m}\left(\prod_{s=1}^{\lambda'_i-\lambda'_{i+1}}(1-p^{-s})\right)\right) p^{\sum_{1\leq i,j\leq m}\min(i,j)(\lambda'_i-\lambda'_{i+1})(\lambda'_j-\lambda'_{j+1})} \\
&= \left(\prod_{i=1}^{m}\left(\prod_{s=1}^{\lambda'_i-\lambda'_{i+1}}(1-p^{-s})\right)\right) p^{\sum_{i=1}^{m}(\lambda'_i)^2}.
\end{aligned}$$

Then, setting $q = p^{-1}$, we have

$$\begin{aligned}
\sum_{G\in\mathcal{G}_p}\frac{1}{|G|^u\cdot|\mathrm{Aut}(G)|} &= \sum_{n=0}^{\infty}\sum_{\substack{G_\lambda\in\mathcal{G}_p\\|\lambda|=n}}\frac{q^{nu}}{|\mathrm{Aut}(G_\lambda)|} \\
&= \sum_{n=0}^{\infty} q^{nu}\sum_{\substack{G_\lambda\in\mathcal{G}_p\\|\lambda|=n}}\left(\prod_{i=1}^{m}\left(\prod_{s=1}^{\lambda'_i-\lambda'_{i+1}}(1-p^{-s})^{-1}\right)\right)\left(\prod_{i=1}^{m}p^{-(\lambda'_i)^2}\right) \\
&= \sum_{n=0}^{\infty} q^{nu}\sum_{\substack{G_\lambda\in\mathcal{G}_p\\|\lambda|=n}}\left(\prod_{i=1}^{m}\left(\prod_{s=1}^{\lambda'_i-\lambda'_{i+1}}(1-q^{s})^{-1}\right)\right)\left(\prod_{i=1}^{m}q^{(\lambda'_i)^2}\right).
\end{aligned}$$

Note that, $\lambda'$ varies over all partitions as $\lambda$ varies over all partitions. Therefore putting $\mu = \lambda'$, we get

$$\sum_{G\in\mathcal{G}_p} \frac{1}{|G|^u \cdot |\mathrm{Aut}(G)|} = \sum_{n=0}^{\infty} q^{nu} \sum_{\substack{G_\mu\in\mathcal{G}_p \\ |\mu|=n}} \left(\prod_{i=1}^{m}\left(\prod_{s=1}^{\mu_i-\mu_{i+1}}(1-q^s)^{-1}\right)\right)\left(\prod_{i=1}^{m} q^{\mu_i^2}\right)$$
$$= \sum_{n=0}^{\infty} q^{nu} \sum_{\substack{G_\mu\in\mathcal{G}_p \\ |\mu|=n}} \left(\prod_{i=1}^{m} \psi_{\mu_i,\mu_{i-1}-\mu_i}(q)\right)\left(\prod_{i=1}^{m} q^{\mu_i^2}\right),$$

where

$$\psi_{a,b}(q) := \frac{\prod_{i=1}^{a+b}(1-q^i)}{\prod_{i=1}^{a}(1-q^i)\prod_{i=1}^{b}(1-q^i)}, \quad \psi_{a,\infty}(q) := \frac{1}{\prod_{i=1}^{a}(1-q^i)}$$

and $\mu_0 := \infty$; note that, the coefficient of $q^n$ in $\psi_{a,b}(q)$ is the number of partitions of $n$ with hight at most $a$ and width at most $b$.Therefore, by Lemma 2, it is enough to show that, for each $n \geq 0$,

$$f_n(q) = \sum_{\substack{G_\mu\in\mathcal{G}_p \\ |\mu|=n}} \left(\prod_{i=1}^{m} \psi_{\mu_i,\,\mu_{i-1}-\mu_i}(q)\right)\left(\prod_{i=1}^{m} q^{\mu_i^2}\right).$$

That is, we need to equate coefficients of $q^N$ on both sides, for each $N \geq 0$. The argument is same as given in [2] or [3].

Note that, coefficient of $q^N$ on LHS is equal to $b_{N,n}$ which is the number of partitions of $N$ with greatest part $n$. Let $\nu$ be a partition of $N$ with greatest part equal to $n$; then, to each such $\nu$ we will associate a partition $\mu$ of size $n$ on the RHS. Consider the conjugate $\nu'$ of $\nu$ and let $D$ be the standard *Young diagram* of $\nu'$. Note that $\nu'$ has hight equal to $n$. Now, define $\mu := (\mu_1, \ldots, \mu_m)$ as follows:

- Define $\mu_1$ to be the largest integer such that $(\mu_1, \mu_1) \in D$.
- For $i \geq 2$, define $\mu_i$ to be the largest integer such that

$$(\mu_1 + \cdots + \mu_i, \mu_i) \in D.$$

(where $(i, j) \in D$ is defined as the block of $D$ situated at the $i$th row from

top and $j$th column from left). Then $|\mu| = n$. If $M$ is the number of blocks outside the squares of size $\mu_i$ then $M = N - \mu_1^2 - \mu_2^2 - \cdots - \mu_m^2$. Let $M_i$ be the number of blocks at the right of the block of size $\mu_i$, i.e.

$$M_i := |\{(x, y) \in D : \mu_1 + \cdots + \mu_{i-1} < x < \mu_1 + \cdots + \mu_i,\ \mu_i < y\}|.$$

Then the blocks corresponding to $M_i$ gives a partition of $M_i$ of hight at most $\mu_i$ and width at most $\mu_{i-1} - \mu_i$ and hence this contributes to the coefficient of $q^{M_i}$ in $\psi_{\mu_i, \mu_{i-1}-\mu_i}(q)$ on RHS. Note that $M = M_1 + \cdots + M_m$ which implies $M_1 + \cdots + M_m + \mu_1^2 + \cdots + \mu_m^2 = N$ and hence $\mu$ contributes to the coefficient of $q^N$ on RHS.

Note that, the above construction can be reversed. Suppose $\mu$ is a partition which corresponds to the coefficient of $q^N$ on RHS such that $\mu$ is specified by the numbers $M_i$, where $M_1 + \cdots + M_m + \mu_1^2 + \cdots + \mu_m^2 = N$, and partitions of $M_i$ of hight at most $\mu_i$ and width at most $\mu_{i-1} - \mu_i$. Then we can construct the Young diagram $D$ and construct the corresponding partition $\nu$ on LHS. Hence, we conclude that the coefficients of $q^N$ on both sides are equal and this proves the theorem.

## References


[1] H. Cohen and H. W. Lenstra, Jr., *Heuristics on class groups of number fields*, Number theory, Noordwijkerhout 1983 (Noordwijkerhout, 1983), Lecture Notes in Math., vol. 1068, Springer, Berlin,1984, pp. 33-62. MR 756082 (85j:11144).

[2] P. Hall, *A partition formula connected with abelian groups*, Comm. Math. Helvetici **11** (1938), no. 1, 126-129.

[3] J. Lengler, *The Cohen-Lenstra Heuristic for Finite Abelian Groups*, PhD thesis, Universität des Saarlandes, 2009.


## Author information


Pritam Majumder, Department of Mathematics and Statistics,
IIT Kanpur, Kanpur-208016, U.P., India.
Email: `pritamaj@gmail.com`